



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

REC'D 24 DEC 2003

WIPO

F

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02080472.0

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr:
Application no.: 02080472.0
Demande no:

Anmeldetag:
Date of filing: 19.12.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

UNILEVER N.V.
Weena 455
3013 AL Rotterdam
PAYS-BAS

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Water continuous product

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

8

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A23L1/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SI SK

Water continuous product

Field of the invention

- 5 The invention relates to water continuous products that are stable against microbiological spoilage.

Background to the invention

- 10 Water continuous products such as oil in water emulsions are prone to microbial spoilage on storage. Among these products are dressings, mayonnaise, fresh cheese, creamy spreads such as those disclosed in WO-A-97/08956 and quark type of products.
- 15 For each of these products measures are taken to ensure that they have a suitable shelf life when closed and also to ensure that they are not spoiled immediately after storage but can safely be consumed for some time after opening. One of the bacteria causing spoilage after the product has been opened are
- 20 those belonging to the family of Enterobacteriaceae.

For microbial preservation many of the above products rely on a preservation system which is a combination of heating (e.g. pasteurisation, sterilisation) and acid level (pH) to kill any

25 vegetative cells present in the product and to inhibit bacterial spores. Heating is in many cases sufficient to ensure closed shelf life prevented products are hermetically sealed after packing. Acid level is e.g. used in dressings where generally the level of acetic acid is about 0.1 to 2 wt% on

30 product weight.

US-A-6387427 for example discloses a method of preserving a food product by adding a fermentate comprising acetic acid and/or its salts resulting from the fermentation by acetic acid producing bacteria while the pH is maintained below 5.8.

5 Although such products may be stable, the production of the fermentate by microbial acidification is time consuming and also the composition of the fermentate may vary depending on the exact conditions of microbial acidification, which are difficult to control very accurately. Furthermore the acetic
10 acid levels are such that for many products an undesired acetic taste and flavour may be present.

Furthermore e.g. products disclosed in WO-A-97/08956 which have been prepared by microbial acidification may be stable but they
15 again require a step wherein microbial acidification is applied which is in some cases undesired. Microbial acidification may take long and makes exact process control of e.g. the pH difficult.

20 Although it is known from the art that microbial acidification may be replaced by chemical acidification, we found that this does not always lead to products with the desired microbial stability. For instance products wherein an acidic pH is obtained by the mere addition of citric acid were found to
25 spoil relatively easily, especially once opened unless they contain very high amounts of citric acid.

Therefore it is an object of the invention to provide chemically acidified products that are microbiologically stable
30 on storage.

Summary of the invention

We suprisingly found that a combination of a food grade acid with a very small amount of acetic acid gives highly improved
5 microbiological stability for water continuous products.

Therefore the invention relates to a water continuous, chemically acidified product comprising a fat, from 0.1 to 15 wt% protein, acids, and having a pH from 3.5 to 6.5 wherein the
10 acids comprise a combination of at least one food grade acid in an amount of from 0.01 to 2 wt% and from 10 to 1000 ppm acetic acid, on the weight of the product.

Detailed description of the invention

15

In the context of the invention, the terms "fat" and "oil" are used interchangeably. The term oil encompasses both triglyceride oils and diglyceride oils. Where the term fat is used also fat replacers such as sucrose polyesters are
20 encompassed.

For the purpose of the current invention, wt% is defined as weight percent on total product weight unless otherwise is indicated.

25

Where the weight amount of an acid is referred to, the amount is calculated in terms of the total amount of dissociated and undissociated acid.

30 The products according to the invention are chemically acidified. This means that the desired acidic pH is obtained by the addition of (food grade) acids, contrary to the other method of acidification which is microbial acidification.

Therefore preferred products are essentially free from bacteria which are suitable for microbial acidification (which is also referred to as fermentation). Examples of such bacteria are lactic acid bacteria, cheese starter cultures, yoghurt

5 cultures, acetic acid producing bacteria. The presence of such bacteria (alive or dead after a heat treatment) may be demonstrated by any suitable technique such as ribosomal RNA identification with fluorescent oligonucleotides or PCR of DNA.

10 The products according to the invention comprise a continuous aqueous phase. The dispersed phase is preferably fat or oil or a replacer thereof. Preferred products are spreadable, liquid and spoonable emulsions such as fresh cheese, spreads, low fat spreads, sauces, creams, drinks, acidified drinks, acidified
15 milk, mousses, desserts, dairy spreads.

The products comprise a protein in an amount from 0.1 to 15 wt%. Especially products with protein present in the aqueous phase were found to be prone to microbial spoilage and hence
20 will benefit from the invention because proteins are a well known source of energy for bacteria to grow on.

The products are acidified to a pH of from 3.5 to 6.5. The water continuous products preferably have a pH from 3.8 to 6.5,
25 more preferably from 4 to 6.0, more preferred from 4.2 to 5.2.

The desired pH is obtained by the addition of a combination of at least one food grade acid in an amount from 0.01 to 2 wt% and from 10 to 1000 ppm acetic acid.

30

The preferred amount of food grade acid is from 0.1 to 1.2 wt%, more preferred from 0.2 to 0.7 wt%. The skilled person will appreciate that the amount of acid added is linked to the

desired pH of the final product and the strength of the particular acid.

It is well known that microbiologically produced acids during fermentation impart a specific, pleasant taste to food products. Therefore in a preferred embodiment, the food grade acid is composed of acids that are formed during fermentation.

The food grade acid is preferably selected from the group comprising citric acid, lactic acid, hydrochloric acid, benzoic acid, propionic acid or a combination thereof.

The most preferred food grade acid is citric acid or a combination of citric acid and lactic acid. In preferred products the total amount of lactic acid is preferably below 0.5 wt%.

The amount of citric acid is preferably from 0.1 to 0.8 wt% , more preferred from 0.2 to 0.4 wt% on total product weight.

It was surprisingly found that the addition of only a small amount of acetic acid showed a very good improvement in inhibition of microbial growth, especially of bacteria of the species Enterobacteriaceae. In a preferred embodiment the amount of acetic acid on total product weight is from 50 to 500 ppm, more preferred from 50 to 300 ppm, even more preferred from 100 to 300 ppm, most preferred from 150 to 250 ppm.

The invention is for example applicable to products which are of a creamy spread type. In such products the level of acetic acid is preferably below 500 ppm to avoid a strong acetic taste for these products. For other products such as dressings the amount of acetic acid may be a bit higher because such products are usually characterised by an acetic acid taste and flavour.

In an even more preferred embodiment, the product comprises citric acid in an amount of from 0.2 to 0.4 wt% and acetic acid in amount from 100 to 300 ppm on total product weight.

5

The products preferably are water continuous products comprising from 1 to 40 wt% vegetable fat, 0.5 to 10 wt% milk protein and 0.01 to 3 wt% thickener, having a pH from 4.2 to 5.8. These products are nutritionally balanced and hence
10 preferably comprise essential nutrients while they are low fat.

The products according to the invention comprise a fat or a fat replacer. Most preferred products are emulsions of a continuous aqueous phase and a dispersed fat phase.

15

The fat or fat replacer is preferably present in an amount of from 5 to 90 wt%, more preferred from 5 to 70 wt%, more preferred from 5 to 50 wt%, even more preferred from 10 to 40 wt%, most preferred from 15 to 35 wt%. The invention is
20 especially beneficial for products with a high amount of aqueous phase such as those where the amount of fat is below 60 wt%. Such products are therefore preferred.

Although any suitable fats such as dairy fat, vegetable fat and
25 fish oil may be used, the preferred fat is vegetable fat. The vegetable fat is preferably a fat rich in polyunsaturated fatty acids. Most preferred the fat is selected from the group comprising palmkernel oil, olive oil, soybean oil, rapeseed oil, coconut oil, sunflower oil, safflower oil, or fully or
30 partially hardened fractions thereof. In a further preferred embodiment, the total amount of saturated fatty acid components in the fat is less than 45 wt%, based on the total amount of

fatty acid components, and further preferred less than about 30 wt%.

The products comprise a protein. The protein may be any protein 5 such as milk protein, soy protein, pea protein, meat protein. Preferred products comprise from 0.5 to 10 wt% protein, preferably from 1.5 to 10 wt%, more preferred from 2 to 5 wt% protein.

- 10 Preferably the protein is milk protein and even more preferred the milk protein is derived from the group comprising cream, milk, skim milk, butter milk, whey, casein or their powders or a combination of any of these. The most preferred milk protein source is skim milk powder.
- 15 The preferred amount of milk protein is from 2 to 6 wt%, more preferred from 2.5 to 4.5 wt%.

Although the invention is applicable for water continuous products of any kind of texture, it is especially preferred 20 that products are spreadable.

Therefore in a preferred embodiment the invention relates to products that are plastic, spreadable products, which can be applied onto bread at room temperature without tearing the bread. The products of the present invention preferably have a 25 Stevens value of 100-500 g at 10°C and 25-250 g at room temperature (about 20°C), more preferred 150-450 g at 10°C and 25-200 g at room temperature (about 20°C). The method to determine Stevens value hardness is described in the examples.

- 30 To obtain the desired consistency and spreadable characteristics, the products according to the invention optionally comprise a thickener. The thickener is preferably a biopolymer selected from the group comprising locust bean gum ,

guar gum, tara gum, amylopectin, methylcellulose, alginate, starch, modified starch, high molecular weight pectin, gelatin or combinations thereof.

The preferred amount of thickener is from 0.1 to 3 wt%, more
5 preferred from 0.3 to 2 wt%.

Optionally the product further comprises an ingredient selected from the group comprising vitamins, anti-oxidants, flavouring agent, egg yolk, gelatin, salt, metal ions such as calcium,
10 potassium or a combination thereof.

Although the acid combination according to the invention has a good contribution to improving the microbial stability of water continuous products against spoilage by especially
15 Enterobacteriaceae, further ingredients are optionally introduced to specifically provide protection against other bacteria such as Listeria. Example of a suitable other anti-microbial agent that is optionally included in the product is potassium sorbate.

20

The products according to the invention are microbiologically stable. The microbial stability may be determined by the test described in the examples.

25 The product may be prepared in any suitable way. A suitable process is for example disclosed in WO-A- 97/08956. It is preferred that the product is filled into a container at a temperature from 60 to 80 °C whereafter the container is stored at a temperature below 10 °C. More preferred the filling

30 temperature is from 65 to 75 °C.

The invention is illustrated in the following non-limiting examples.

Examples

5

Stevens hardness

The firmness of the products is determined by measuring the force required to penetrate a cylindrical probe in the product. Sample height 5 cm; cylindrical probe of 0.5 inch thickness; 10 compression rate 2 mm/s; penetration depth 20 mm. The samples are stored for 7 days at 5 °C, and stored at 10 or 20°C for 4 h before the firmness measurement.

Microbial stability test

15

The microbiological stability of the products is determined using a "challenge" test which is a common test in the determination of microbiological stability.

The test was carried out as follows.

20 Products were opened and a cocktail of strains of the Enterobacteriaceae family was added. Products were incubated at 10, 15 and 25 °C for several days. The amount of growth of Enterobacteriaceae was determined by plating decimal dilution series of the "infected" products on agar plates. The plates 25 were incubated at 30 °C for 2 to 5 days after which the numbers of grown colonies were counted. Typically for Enterobacteriaceae the increase of the number of colonies was determined versus time.

Example 1

Table 1: Compositions

Ingredient	Wt % on product
Fat (sunflower oil)	22
Skim Milk Powder	6.25
Whey protein concentrate (Nutrilac)	1.75
Locust Bean Gum (LBG)	0.3
Gelatin	0.7
Salt	0.3
Citric acid (added as a powder)	0.35
Acetic acid (added as 100% liquid)	200 ppm
Carotene stock solution, 1% in oil	4 ppm
Demineralised water	Up to 100%

Comparative example C1: same ingredients except that there was no acetic acid.

Process:

5

Water phase and fat phase ingredients except for acids were mixed at about 60 °C. After mixing the composition was pasteurized at 85°C for 10 minutes, and cooled down to 44°C, after which homogenisation at 200 bar took place. To the
10 homogenized composition acid was added as indicated, where needed hydrochloric acid was added until a pH of about 4.7 or 4.5 was reached. This was followed by heating the mixture to 85 °C. The obtained product was homogenized at 300 bar, and subsequently heated to a temperature of 75°C for filling
15 containers. The product was filled into the containers. The container was stored below 10°C.

The product was presented to a panel of consumers. The products were easily spreadable. All confirmed that both products had a
20 pleasant mouthfeel and taste and did not have an acetic acid off taste.

The results of the microbial stability tests were:

The samples with acetic acid (according to the invention) did
25 not show any increase of colonies of Enterobacteriaceae over time. After about 10 days at 10 °C, even a reduction in the initial number of colonies was observed at pH 4.7.

For products at pH 4.7 without acetic acid (comparative
30 example) growth of Enterobacteriaceae started between about 3 to 7 days storage at 10 °C. After about 10 days the amount of colonies had increased about 10-fold.

For products at pH 4.5 without acetic acid, the growth of colonies was deferred but was observed to start after storage for 10 to 14 days at 10 °C.

5

This confirms that the addition of a small amount of acetic acid inhibits microbial growth of Enterobacteriaceae. Therefore such products have a longer open shelf life.

EPO - DG 1
19 12 2002

13

(44)

Claims

1. Water continuous, chemically acidified product comprising a fat, from 0.1 to 15 wt% protein, acids, and having a pH from 3.5 to 6.5 characterised in that the acids comprise a combination of at least one food grade acid in an amount of from 0.01 to 2 wt% and from 10 to 1000 ppm acetic acid, on the weight of the product.
2. Water continuous product according to claim 1 wherein the food grade acid is selected from the group comprising citric acid, lactic acid, hydrochloric acid, benzoic acid, propionic acid or a combination thereof.
3. Water continuous product according to claim 2 wherein the food grade acid is citric acid or a combination of citric acid and lactic acid.
4. Water continuous product according to any of claims 1-3 with a pH from 3.8 to 6.5, preferably from 4 to 6.0, more preferred from 4.2 to 5.2.
5. Water continuous product according to any of claims 1-4, which comprises from 5 to 90 wt% oil.
6. Water continuous product according to claim 1 or 2 which comprises citric acid in an amount of from 0.2 to 0.4 wt% and acetic acid in amount from 100 to 300 ppm on total product weight.
7. Water continuous product according to claim 1 wherein the amount of fat is from 1 to 40 wt% fat, said product further

comprising 0.5 to 10 wt% milk protein and 0.01 to 3 wt% thickener and having a pH from 4.2 to 5.8.

8. Water continuous product according to any of claims 1-7 wherein the amount of lactic acid is below 0.5 wt% on total product weight.

Abstract

Chemically acidified water continuous products which have been acidified with a combination of a food grade acid and a small amount of acetic acid were found to be very stable against microbial spoilage.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.